

POTENTIAL OF PEDADA FRUIT AS A RENEWABLE ENERGY AND CLIMATE MITIGATION SOURCE IN INDONESIA: A REVIEW

Sari Sekar Ningrum^{1*}, Annisa Yuliana Angeline², Muhamad Sofi Ardani², Zhilal Shadiq¹

¹⁾ Teknologi Rekayasa Kimia Industri, Politeknik Negeri Cilacap, Indonesia

²⁾ Teknik Pengendalian Pencemaran Lingkungan, Politeknik Negeri Cilacap, Indonesia

Abstract

Indonesia has one of the largest mangrove ecosystems in the world, which provides important ecological functions such as coastal protection, biodiversity support, and climate mitigation. Among various mangrove species, *Sonneratia* sp. (pedada fruit) has attracted attention due to its high carbohydrate content, making it a promising biomass source for renewable energy production. This study aims to evaluate the potential of pedada fruit as a raw material for bioethanol production and its contribution to climate change mitigation in Indonesia through a comprehensive literature review. This review analyzed approximately 50 scientific articles, conference proceedings, and research reports related to mangrove biomass utilization, bioethanol production processes, and the environmental benefits of mangrove ecosystems. The results showed that pedada fruit contains approximately 59% carbohydrates along with lignocellulosic components such as cellulose, lignin, and hemicellulose, which are suitable for conversion into bioethanol through pretreatment, hydrolysis, fermentation, and distillation processes, and can be applied in Indonesia with its large number of mangroves. Bioethanol production from mangrove fruit can follow second- to fourth-generation bioethanol technologies, depending on the processing approach used. In addition to providing renewable energy, utilizing mangrove fruit as a bioethanol feedstock can encourage community participation in mangrove conservation and restoration efforts. Therefore, mangrove fruit has significant potential as a sustainable, renewable energy source while also supporting climate mitigation strategies and coastal ecosystem management in Indonesia.

Keywords: *bioethanol, climate mitigation, mangrove, renewable energy, Sonneratia sp.*

Introduction

Indonesia is the largest archipelagic country in the world, with a sea area of more than 70% of the total area of Indonesia (Rencana Strategis Deputy Bidang Koordinasi Kedaulatan Maritim Dan Energi Tahun 2020-2024, 2020). Coastal areas in Indonesia are widely overgrown with mangrove plants. Mangroves function as

wave breakers, prevent abrasion, produce food for living things, and in climate mitigation efforts (Dinilhuda et al., 2019) so that mangrove plants are very important plants to be planted in Indonesia. The absence of mangroves can cause a lack of natural protection for the coastline, so that coastal ecosystems become more easily damaged by human activities and extreme weather changes (Irfandi et al., 2025). Based on data from the Ministry of Forestry in 2025, the area of mangroves in Indonesia is 3,440,464 hectares for existing ones and 769,824 hectares for the area of potential mangrove habitat (Keputusan Menteri

*Corresponding Author:

E-mail: sarisekarningrum@pnc.ac.id

Received: 13 March 2026

Revised: 18 March 2026

Accepted: 19 March 2026

DOI: 10.23969/jcbeem.v10i1.43632

Kehutanan Republik Indonesia No 594 Tahun 2025 Tentang Peta Mangrove Nasional 2024, 2025). Mangroves in Indonesia have a lot of species diversity, including *Rhizophora* sp., *Avicenna* sp., *Sonneratia* sp., *Bruguire* sp., *Xylocarpus* sp., *Ceriops* sp., and *Exoecaria* sp. (Fitria & Dwiyanoto, 2021). These various types of mangroves contain carbohydrates (Fattah et al., 2025), so they can be used to make further products, one of which is bioethanol, a renewable energy source that can replace the use of fossil fuels.

With this background, a literature review was conducted to evaluate the potential of pedada fruit (*Sonneratia* sp.) to become bioethanol as renewable energy in Indonesia, with the hope of providing a comprehensive picture to strengthen the role of pedada fruit as bioethanol in the national energy transition.

Based on research conducted by (Huda, 2017), raw materials that are considered efficient and easy to process into bioethanol generally have a carbohydrate content of between 50-60% of the dry weight. Of the various types of mangroves that have a carbohydrate content of between 50-60% is *Sonneratia* sp. *Sonneratia* sp or better known as pedada fruit, is one type of mangrove fruit that has a carbohydrate content of between 50-60%. From research conducted by (Mila Prametha et al., 2025), the carbohydrate content in pedada fruit is around 59.09%, so it has the potential to produce bioenergy, especially bioethanol. In addition to the carbohydrate content, the content of lignin, cellulose, and hemicellulose can also be processed into bioethanol.

Energy consumption in Indonesia increases annually, driven by the increasing number of industries, transportation, households, and agriculture. According to data from the Central Statistics Agency (BPS), energy consumption increased by 6.4% in 2023 compared to 2022. Fossil fuels account for approximately 90% of the country's primary energy source (Kementerian PPN/Bappenas, 2024). The use of fossil fuels can lead to greenhouse gas emissions, which can contribute to global warming, climate change, air pollution, and environmental damage (Anser et al., 2020; Breetz et al., 2025; Li, 2023; Steel et al., 2025; Zimon et al., 2023). Therefore, a transition to a sustainable energy system based on renewable energy is needed.

Bioethanol is a renewable energy source that can replace fossil fuels (Tobe et al., 2022). Bioethanol plays a crucial role in the global transition to sustainability, serving as a renewable fuel, particularly in the transportation sector, and reducing greenhouse gas emissions (Vacharanukrauh et al., 2025). Bioethanol is a renewable energy source derived from carbohydrate-containing raw materials through a fermentation process (Kahar et al., 2025).

Research Methodology

This study employed a comprehensive literature review method involving extensive and in-depth research on the potential of pedada fruit as a bioethanol feedstock. The information sources were extensive and varied, including research reports, conference proceedings, and relevant scientific articles. The research articles used as references in this study were taken from the last 10 years of data. The stages of the

literature review method include: 1) collecting research literature related to bioethanol, 2) analyzing and synthesizing data, 3) interpreting data, and 4) concluding. The data in this study were collected from various secondary sources derived from research articles focusing on the same research topic.

Results and Discussion

Potential of Pedada Fruit

Pedada fruit (*Sonneratia* sp.) is a plant that lives in the mangrove ecosystem (Handayani et al., 2025). Pedada fruit can be converted into several energy products, as shown in Figure 1.




Figure 1. Various energy products produced by pedada fruit

Pedada fruit contains carbohydrates with a content of 59.09%, which has a function as an energy producer (Mila Prametha et al., 2025). In addition to containing carbohydrates, pedada fruit also contains 24.8995% cellulose, 5.0896% lignin, and 11.7486% hemicellulose (Ningrum et al., 2025). The content of carbohydrates, cellulose, lignin, and hemicellulose in pedada fruit can be converted into bioethanol (Beluhan et al., 2023; Damayanti et al., 2021; Darsono & Sumarti, 2014; Dwicahyana et al., 2025).

Development of Bioethanol Production Process

Currently, bioethanol production is divided into four generations based on the raw materials used in the process of producing bioethanol (Mohd Azhar et al., 2017; Tse, Wiens, & Reaney, 2021). In all generations, through the process of pre-treatment, hydrolysis, fermentation, and separation/purification (Vacharanukrauh et al., 2025). The pre-treatment process is very important in the process of making bioethanol in some raw materials because it affects the release of sugar in the fermentation process, which can affect the final result of bioethanol (Ben-Iwo et al., 2016; Tse, Wiens, Shen, et al., 2021).

The development of the bioethanol production process began with the first generation, which utilized raw materials with high starch content, such as corn and wheat seeds, and sugar sources like sugar cane, thereby causing competition between food and energy (Petroleum Supply Annual, 2024; Puri et al., 2012; Rebeiro, 2013). In the second generation, bioethanol is developed using raw materials that are not sourced as food ingredients, namely, using raw materials sourced from lignocellulose, which is abundant and sustainable (Zuliani et al., 2021). In the third generation, bioethanol is produced using microorganisms that synthesize fermentable carbohydrates, making the process possible through enzymatic hydrolysis and microbial fermentation (Dutta et al., 2014; Nishida et al., 2023). Fourth-generation bioethanol is the most advanced stage in bioethanol development, utilizing metabolic engineering and synthetic biology to

increase fermentation efficiency (Assaf et al., 2024; Mignogna et al., 2024). In this fourth generation, several microorganisms are genetically engineered to utilize glucose and pentose sugars found in lignocellulosic biomass hydrolysates for bioethanol production.

Bioethanol Production with Raw Materials from Pedada Fruit

Pedada fruit is a type of mangrove that contains carbohydrates, cellulose, lignin, and hemicellulose. Pedada fruit can be used as raw material for bioethanol. The process of producing bioethanol can follow the process in the second, third, or fourth generation if seen from the content contained in the pedada fruit.

Initial Treatment

In the manufacture of bioethanol, pretreatment is very crucial. This pretreatment aims to change the physical and chemical structure of biomass so that the sugars in it are more easily converted into bioethanol (Humera Farheen et al., 2026; Khairiah & Ridwan, 2021; Shukla et al., 2023). This treatment process includes cleaning and drying to remove dirt and reduce the water content in the raw material so that the raw material is ready to be processed in the fermentation stage (Dayatmo & HS, 2015; Jasman & Ahmad, 2021; Jönsson & Martín, 2016).

Fermentation Process

The content parameters in pedada fruit are a very important part of determining the fermentation route that will be carried out. The content of pedada fruit can be seen in Table 1.

Table 1. Pedada fruit content (Ningrum et al., 2025)

Pedada Fruit Content	Amount (%)
Water content	10.8399
Ash content	5.8708
Lipid content	0.6123
Protein content	8.1004
Carbohydrate content	59.0948

Table 1 shows that pedada fruit (*Sonneratia* sp.) has a low protein content but a high carbohydrate content. This high carbohydrate content makes it suitable as a raw material for bioethanol because carbohydrates can be fermented directly into bioethanol (Amalia & Rezeki Muria, n.d.; Andana et al., 2020; Sadimo et al., 2016). Fermentation of carbohydrates into bioethanol can use several microorganisms, including *Saccharomyces cerevisiae*, *Zymomonas mobilis*, *Trichoderma viride*, and several types of indigenous yeast (Gustina et al., 2022; Mohd Azhar et al., 2017; Praveen & Brogi, 2025). This fermentation process is carried out to convert the reducing sugar content obtained from carbohydrates in pedada fruit into bioethanol.

Distillation Process

The distillation process is one of the techniques used in bioethanol purification (Onuki et al., 2008). The distillation process is carried out by heating a liquid mixture, where the components with the lower boiling points will evaporate first, concentrate in the vapor phase, and then condense back into a liquid (Setiawan, 2018; Stewart, 2024). This distillation method is the most common method used to increase the alcohol content of fermentation products

in the production of bioethanol (Lay et al., 2010; Sehwantoro et al., 2021).

Mangrove Plants as a Form of Climate Mitigation Effort in Indonesia

Mangroves in Indonesia play a vital role in climate mitigation. Mangroves have the capacity to absorb 3-5 times more carbon than tropical forests and can protect coastal areas from abrasion, tsunamis, and sea level rise (Fitria & Dwiyanoto, 2021; Soleman Imburi et al., 2024; Wihartono, 2025). The use of mangrove fruit as bioethanol can increase community motivation to plant and care for mangroves (Abubakar et al., 2023; Anggraeni Luthfiyatul Afifah et al., 2025; Farhaeni, 2016; Mulyani et al., 2018).

Conclusions

Based on the nutritional content of pedada fruit (*Sonneratia* sp.), it can be concluded that pedada fruit has potential as a raw material for bioethanol, a renewable energy source. This content allows pedada fruit to be processed into bioethanol through several stages, including pre-treatment, hydrolysis, fermentation, and distillation. Based on a literature review, bioethanol production from pedada fruit can be integrated with second-, third-, or even fourth-generation bioethanol technologies.

In addition to its potential as a bioenergy source, the utilization of pedada fruit for bioethanol production can contribute to climate change mitigation and sustainable coastal management. By providing economic value to mangrove resources, this approach may encourage local communities to participate in mangrove conservation and restoration efforts. Therefore, the development of pedada-based bioethanol

can support Indonesia's transition toward renewable energy while simultaneously strengthening ecosystem protection and community-based environmental management.

Acknowledgment

The author would like to express his gratitude to the Politeknik Negeri Cilacap which has always provided support to the author in conducting research.

References

- Abubakar, S., Abdul Kadir, M., Subur, R., Fadel, A. H., Said Al Hadad, M., Wahidin, N., Noman Susanto, A., Dewi Salim, F., Muksin, D., haddad, A., & N, S. A. (2023). Pemanfaatan Buah Mangrove *Rhizophora apiculata* Sebagai Olahan Kopi Mangrove Dalam Upaya Peningkatan Ekonomi Masyarakat di Desa Maitara Utara Kecamatan Tidore Utara. *Jurnal Pengabdian Magister Pendidikan IPA*, 6(2), 368–377. <https://doi.org/10.29303/jpmpi.v6i2.4476>
- Amalia, Y., & Rezeki Muria, S. (n.d.). *Pembuatan Bioetanol dari Limbah Padat Sagu Menggunakan Enzim Selulase dan Yeast Saccharomyces Cerevisiae dengan Proses Simultaneous Sacharification and Fermentation (SSF) dengan Variasi Konsentrasi Substrat dan Volume Inokulum*.
- Andana, A. D., Tjahjani, S., & Amaria, D. (2020). Penggunaan Antioksidan Sebagai Upaya untuk Menghambat Proses Oksidasi Bioetanol dari Singkong Karet (*Manihot glaziovii*) The Use of Antioxidant to Increase Bioethanol Oxidation Process from

- Rubber Cassava (*Manihot glaziovii*). *UNESA Journal of Chemistry* (Vol. 9, Number 1).
- Anggraeni Luthfiyatul Afifah, As Dewi Aman Meker, Wedar Putri Sholehati, Nur Elsa Choiru Ummah, & Muhammad Yusron Maulana El-Yunusi. (2025). Kegiatan Penanaman Kembali Bibit Mangrove untuk Pelestarian Buah Bakau sebagai Bahan Pembuatan Sirup di Desa Wonorejo Surabaya. *Manfaat : Jurnal Pengabdian Pada Masyarakat Indonesia*, 2(2), 01–16. <https://doi.org/10.62951/manfaat.v2i2.313>
- Anser, M. K., Hanif, I., Alharthi, M., & Chaudhry, I. S. (2020). Impact of fossil fuels, renewable energy consumption and industrial growth on carbon emissions in Latin American and Caribbean economies. *Atmosfera*, 33(3), 201–213. <https://doi.org/10.20937/ATM.52732>
- Assaf, J. C., Mortada, Z., Rezzoug, S. A., Maache-Rezzoug, Z., Debs, E., & Louka, N. (2024). Comparative Review on the Production and Purification of Bioethanol from Biomass: A Focus on Corn. *Processes*, 12(5), 1001–1025. <https://doi.org/10.3390/pr12051001>
- Beluhan, S., Mihajlovski, K., Šantek, B., & Ivančić Šantek, M. (2023). The Production of Bioethanol from Lignocellulosic Biomass: Pretreatment Methods, Fermentation, and Downstream Processing. *Energies*, 16(19), 7003–7042. <https://doi.org/10.3390/en16197003>
- Ben-Iwo, J., Manovic, V., & Longhurst, P. (2016). Biomass resources and biofuels potential for the production of transportation fuels in Nigeria. *Renewable and Sustainable Energy Reviews*, 63, 172–192. <https://doi.org/10.1016/j.rser.2016.05.050>
- Breetz, H. L., Shelton, R., Kaul, M., & Kunkel, L. C. (2025). Just transition funds in U.S. states: explaining variation in political process and policy design. *Environmental Research: Energy*, 2(4), 1–25. <https://doi.org/10.1088/2753-3751/add93c>
- Damayanti, D., Supriyadi, D., Amelia, D., Saputri, D. R., Devi, Y. L. L., Auriyani, W. A., & Wu, H. S. (2021). Conversion of lignocellulose for bioethanol production, applied in bio-polyethylene terephthalate. *Polymers*, 13(17), 2886–2916. <https://doi.org/10.3390/polym13172886>
- Darsono, & Sumarti, M. (2014). Pembuatan Bioetanol dari Lignoselulosa Tandan Kosong Kelapa Sawit Menggunakan Perlakuan Awal Iradiasi Berkas Elektron dan NaOH. *J. Kimia Kemasan*, 36(2), 245–253. <https://doi.org/10.24817/jkk.v36i2.1891>
- Dayatmo, D., & HS, H. (2015). Pembuatan Bioethanol dari Limbah Ampas Pati Aren dengan Metode Hidrolisis Enzimatis menggunakan Enzim Ligninolitik dari Jamur Pelapuk Putih. *Jurnal Konversi*, 4(2), 43-52.
- Dinilhuda, A., Akbar, A. A., & Jumiati. (2019). Peran Ekosistem Mangrove Bagi Mitigasi Pemanasan Global. *Journal of Civil Engineering, University of Tanjungpura*, 18, 191–

198.
<https://doi.org/https://doi.org/10.26418/jtst.v18i2.31233>
- Dutta, K., Daverey, A., & Lin, J.-G. (2014). Evolution Retrospective for Alternative Fuels: First to Fourth Generation. *Renewable Energy*, 69, 114–122.
<https://doi.org/https://doi.org/10.1016/j.renene.2014.02.044>
- Dwicahyana, K. A., Kusuma, I. G. B. W., & Lokantara, I. P. (2025). Proses Pembuatan Bioetanol Dari Buah Mangrove. *Jurnal Ilmiah Teknik Desain Mekanika*, 14(3).
- Farhaeni, M. (2016). Komodifikasi Ragam Buah Mangrove untuk Pemberdayaan Masyarakat Pesisir di Desa Tuban, Kecamatan Kuta, Kabupaten Badung Bali. *Jurnal Studi Kultural*, 1(1), 21–27.
<http://journals.an1image.net/index.php/ajsk>
- Fattah, M., Hakim, L., Soemarno, S., & Purwanti, P. (2025). Analisis pemanfaatan buah mangrove *Rhizophora mucronata* dalam mendukung wisata mangrove masyarakat Kampung Mandar Kabupaten Banyuwangi. *Jurnal Sumberdaya Akuatik Indopasifik*, 9(2), 87–98. <https://doi.org/10.46252/jsai-fpik-unipa.2025.vol.9.no.2.518>
- Fitria, A., & Dwiyanto, G. (2021). Jurnal Ekologi, Masyarakat & Sains Ekosistem Mangrove dan Mitigasi Pemanasan Global. *Jurnal Ekologi, Masyarakat & Sains*, 2(1), 2021. <http://journals.ecotas.org/index.php/e ms>
- Gustina, M., Jalaluddin, ZA, N., Bahri, S., & Masrullita. (2022). Pengaruh Lama Waktu Fermentasi Terhadap Kadar Bioetanol Dari Pati Ubi Jalar Ungu (*Ipomea batata* L). *Chemical Engineering Journal Storage*, 2(2), 116–125.
- Handayani, R., Pagarra, H., Mu'nisa, A., & Hala, Y. (2025). Uji Pengaruh Ekstrak Etanol Buah Pedada (*Sonneratia caseolaris*) Terhadap Penurunan Kadar Asam Urat Mencit (*Mus musculus*) Jantan. *BioEksakta: Jurnal Ilmiah Biologi Unsoed*, 7(3), 265–272. <https://doi.org/10.20884/1.bioe.2025.7.3.17337>
- Huda, N. (2017). *Proses Pembuatan Bioethanol*.
- Humera Farheen, V., Aslam Abdullah, M., & Ganesh Moorthy, I. (2026). Lignocellulosic bioethanol production: a review on pretreatment strategies, biofuel separation, and artificial intelligence/machine learning – based sustainable optimization. *Current Research in Biotechnology*, 11. <https://doi.org/10.1016/j.crbiot.2025.100355>
- Irfandi, M., Nur Mualimin, C., Rahayu Sianturi, P., Aura Magfira, N., Riski Mulya, U., Kholyfaur, U. R., Deri, L., Hidayat Ardiansyah Hady, M., Perayana, Amelia Rossa, S., & Asni. (2025). Penanaman Pohon Mangrove Sebagai Upaya Pelestarian Lingkungan Di Desa Konaweha. *Jurnal Abdimas Indonesia*, 5(4), 2603–2611. <https://dmi-journals.org/jai/>
- Jasman, & Ahmad, R. M. (2021). Pengaruh Jenis Perlakuan Awal terhadap Konsentrasi Bioetanol Hasil Hidrolisis dan Fermentasi Tongkol Jagung menggunakan *Trichoderma reesei* dan

- Saccharomyces cerevisiae*. *Jurnal Beta Kimia*, 1(2). <http://ejurnal.undana.ac.id/index.php/jbkHalaman|25>
- Jönsson, L. J., & Martín, C. (2016). Pretreatment of lignocellulose: Formation of inhibitory by-products and strategies for minimizing their effects. *Bioresource Technology*, 199, 103–112. <https://doi.org/10.1016/j.biortech.2015.10.009>
- Kahar, A., Fathoni, an, Ardiah, R., & Mulyani, N. (2025). Proses Pembuatan dan Uji Karakteristik Bioetanol dari Kulit Nanas (Ananas Comosus). *Jurnal Chemurgy*, 9(1), 58–64. <https://doi.org/http://dx.doi.org/10.30872/cmg.v9i1.16350>
- Kementerian PPN/Bappenas. (2024). *Energi*. Kementerian PPN/Bappenas. <https://lcdi-indonesia.id/grk-energi#:~:text=Bagi%20Indonesia%20yang%20memiliki%20proporsi,untuk%20melakukan%20dekarbonisasi%20semakin%20tinggi>
- Keputusan Menteri Kehutanan Republik Indonesia No 594 Tahun 2025 Tentang Peta Mangrove Nasional 2024 (2025).
- Khairiah, H., & Ridwan, M. (2021). Pengembangan Proses Pembuatan Bioetanol Generasi II dari Limbah Tandan Kosong Kelapa Sawit. *Jurnal Pangan Dan Agroindustri*, 9(4), 233–240.
- Lay, A., Pasang, P. M., & Iqbal, T. A. (2010). Destilasi-Dehidrasi Bioetanol dari Nira Aren dan Karakteristiknya. *Buletin Pama*, 39.
- Li, X. (2023). How can we make an orderly transition away from fossil fuels? A global tour of modern energy. *Sage Journals*, 35(6), 2959–2974. <https://doi.org/https://doi.org/10.1177/0958305X231161300>
- Mignogna, D., Szabó, M., Ceci, P., & Avino, P. (2024). Biomass Energy and Biofuels: Perspective, Potentials, and Challenges in the Energy Transition. *Sustainability (Switzerland)*, 16(16), 7036–7069. <https://doi.org/10.3390/su16167036>
- Mila Prametha, N., Aza Fauziana, N., Yuliana Angelin, A., Sofi Ardani, M., & Sari Sekar Ningrum, dan. (2025). Pengaruh Waktu Terhadap Kandungan Antioksidan Dan Vitamin C Dari Ekstrak Buah Pedada (Sonneratia Caseolaris) Sebagai Bahan Baku Sediaan Lotion. *Jurnal Teknologi*, 13(1), 2025–2050. <https://doi.org/10.31479/jtek.v13i1.425>
- Mohd Azhar, S. H., Abdulla, R., Jambo, S. A., Marbawi, H., Gansau, J. A., Mohd Faik, A. A., & Rodrigues, K. F. (2017). Yeasts in sustainable bioethanol production: A review. *Biochemistry and Biophysics Reports*, 10, 52–61. <https://doi.org/10.1016/j.bbrep.2017.03.003>
- Mulyani, Y., Lewaru, M. W., & Haetami, K. (2018). Pemanfaatan dan Pelestarian Mangrove untuk Meningkatkan Kesejahteraan Masyarakat Pesisir Pangandaran. *Jurnal Pengabdian Kepada Masyarakat*, 11(2).
- Ningrum, S. S., Ardani, M. S., Angeline, A. Y., Fauziana, N. A., & Prametha, N. M. (2025). *Ekstraksi Antioksidan dari Buah Pedada sebagai Bahan Pembuat Lotion*.

- Nishida, V. S., Woiciechowski, A. L., Valladares-Diestra, K. K., Zevallos Torres, L. A., Vandenberghe, L. P. de S., Zandoná Filho, A., & Soccol, C. R. (2023). Second Generation Bioethanol Production from Soybean Hulls Pretreated with Imidazole as a New Solvent. *Fermentation*, *9*(2), 93–106. <https://doi.org/10.3390/fermentation9020093>
- Onuki, S., Koziel, J. A., Van Leeuwen, H., Jenks, W. S., Grewell, D., & Cai, L. (2008). Ethanol production, purification, and analysis techniques: a review. *An ASABE Meeting Presentation*.
- Petroleum Supply Annual. (2024, February). *Biofuels explained*. <https://www.eia.gov/energyexplained/biofuels/>
- Praveen, M., & Brogi, S. (2025). Microbial Fermentation in Food and Beverage Industries: Innovations, Challenges, and Opportunities. *Foods*, *14*(1). <https://doi.org/10.3390/foods14010114>
- Puri, M., Abraham, R. E., & Barrow, C. J. (2012). Biofuel production: Prospects, challenges and feedstock in Australia. *Renewable and Sustainable Energy Reviews*, *16*(8), 6022–6031. <https://doi.org/https://doi.org/10.1016/j.rser.2012.06.025>
- Rebeiro, B. E. (2013). Beyond commonplace biofuels: Social aspects of ethanol. *Energy Policy*, *57*, 355–362. <https://doi.org/https://doi.org/10.1016/j.enpol.2013.02.004>
- Rencana Strategis Deputi Bidang Koordinasi Kedaulatan Maritim dan Energi Tahun 2020-2024, Pub. L. 169/Di Tahun 2020 (2020). <https://maritim.go.id/uploads/magazine/20230712090137-2023-07-12magazine090134.pdf>
- Sadimo, M. M., Said, I., & Mustapa, K. (2016). *Pembuatan Bioetanol dari Pati Umbi Talas (Colocasia esculenta L Schott) melalui Hidrolisis Asam dan Fermentasi*. *5* (2).
- Sehwantoro, W., Hindarti, F., & Oktivina, M. (2021). Rancang Bangun dan Uji Kinerja Destilator Elektrik Sebagai Alat Destilasi pada Proses Pembuatan Bioethanol. *Sainstech Jurnal Penelitian dan Pengkajian Sains Dan Teknologi*, *31*(2), 1–10. <https://doi.org/10.37277/stch.v31i2.1125>
- Setiawan, T. (2018). Rancang Bangun Alat Destilasi Uap Bioetanol dengan Bahan Baku Batang Pisang. *Jurnal Media Teknologi*, *04*(02).
- Shukla, A., Kumar, D., Girdhar, M., Kumar, A., Goyal, A., Malik, T., & Mohan, A. (2023). Strategies of pretreatment of feedstocks for optimized bioethanol production: distinct and integrated approaches. *Biotechnology for Biofuels and Bioproducts*, *16*(1). <https://doi.org/10.1186/s13068-023-02295-2>
- Soleman Imburi, C., Angrianto, R., Tanur, E. A., Widodo, I., & Sitompul, G. A. (2024). Peran Hutan Mangrove dalam Menanggulangi Dampak Perubahan Iklim di Wilayah Pesisir Indonesia. *Jurnal Geosains West Science*, *2*(03), 122–132. <https://doi.org/10.58812/jgws.v2i03.1678>
- Steel, D., Bartha, P., & Cripps, R. (2025). Climate Precaution and Producer

- versus Consumer Dependence on Fossil Fuels. *Ethics, Policy and Environment*, 28(1), 1–25. <https://doi.org/10.1080/21550085.2023.2247815>
- Stewart, M. V. (2024). *Distillation*. EBSCO. <https://www.ebsco.com/research-starters/chemistry/distillation>
- Tobe, A. Y., Riwu, D. B. N., Pah, J. C. A., Adoe, D. G. H., & Baitanu, R. (2022). Temperatur, Warna Nyala Api, dan Tinggi Api pada Pembakaran Premixed Bioetanol dari Lontar (*Borassus Flabellifer*). *Lontar*, 9(1), 84–89. <https://doi.org/https://doi.org/10.35508/ljtmu.v9i01.8062>
- Tse, T. J., Wiens, D. J., & Reaney, M. J. T. (2021). Production of Bioethanol—A Review of Factors Affecting Ethanol Yield. *Fermentation*, 7(4), 268–286. <https://doi.org/10.3390/fermentation7040268>
- Tse, T. J., Wiens, D. J., Shen, J., Beattie, A. D., & Reaney, M. J. T. (2021). *Saccharomyces cerevisiae* Fermentation of 28 Barley and 12 Oat Cultivars. *Fermentation*, 7(2), 59–71. <https://doi.org/10.3390/fermentation7020059>
- Vacharanukrauh, T., Soottitantawat, A., Thongchul, N., Kiatkittipong, W., Weeranoppanant, N., & Assabumrungrat, S. (2025). A review on comprehensive strategies for decarbonizing bioethanol production process. *Green Energy and Resources*, 3(4), 100153. <https://doi.org/10.1016/j.gerr.2025.100153>
- Wihartono, A. Y. (2025, December). *Mitigasi Bencana Perubahan Iklim, Pemprov Bangka Belitung dan YKAN Konsultasikan Rencana Aksi Konservasi dan Pemulihan Ekosistem Mangrove*. Yayasan Konservasi Alam Nusantara. <https://www.ykan.or.id/id/publikasi/artikel/siaran-pers/mitigasi-bencana-perubahan-iklim-rencana-aksi-konservasi-dan-pemulihan-ekosistem-mangrove/#:~:text=Manajer%20Program%20Karbon%20Biru%20YKAN,penghidupan%20masyarakat%20pesisir%20Bangka%20Belitung>
- Zimon, G., Pattak, D. C., Voumik, L. C., Akter, S., Kaya, F., Walasek, R., & Kochański, K. (2023). The Impact of Fossil Fuels, Renewable Energy, and Nuclear Energy on South Korea's Environment Based on the STIRPAT Model: ARDL, FMOLS, and CCR Approaches. *Energies*, 16(17), 1–21. <https://doi.org/10.3390/en16176198>
- Zuliani, L., Serpico, A., De Simone, M., Frison, N., & Fusco, S. (2021). Biorefinery gets hot: Thermophilic enzymes and microorganisms for second-generation bioethanol production. *Processes*, 9(9), 1583–1601. <https://doi.org/10.3390/pr9091583>